



CELL LAB Hamster Anti-Mouse CD28

<u>Cat. No.</u>	<u>Form</u>	<u>Quantity</u>
732100	Purified (UNLB) Antibody	0.5 mg
732101	Fluorescein (FITC) Conjugate	0.5 mg
732102	Biotin (BIOT) Conjugate	0.5 mg
732103	Phycoerythrin (PE) Conjugate	0.1 mg
733284	Phycoerythrin (PE) Conjugate	0.2 mg
732104	Allophycocyanin (APC) Conjugate	0.1 mg
732105	Spectral Red™ (SPRD) Conjugate	0.1 mg

For Laboratory Use Only

DESCRIPTION

Clone: 37.51
Isotype: Syrian Hamster IgG λ
Specificity: CD28 co-stimulatory receptor, Mr 40 kDa

CD28 is a type I disulfide-linked homodimer that is constitutively expressed on most thymocytes, at low density on nearly all CD4⁺ and CD8⁺ peripheral T lymphocytes, and at very low levels on NK cells. Its expression is upregulated upon T-cell activation.¹⁻⁴ CD28 is a ligand for CD80/B7-1 and CD86/B7-2 on B cells and other antigen presenting cells,⁵ and plays an important role in the interaction between T cells and B cells. CD28 is a co-stimulatory receptor involved in many, but not all, T-cell independent immune responses.⁵⁻⁸

APPLICATIONS

- Flow cytometry²⁻⁵
- Immunoprecipitation²
- *In vitro* co-stimulation of T cells and NK cells^{1,4,7}

CHARACTERIZATION

To ensure lot-to-lot consistency, each batch of product is tested to conform with characteristics of a standard reference reagent using immunofluorescence staining and flow cytometry.

WORKING DILUTIONS

Flow Cytometry:

FITC conjugate	≤2 μg/10 ⁶ cells
BIOT conjugate	≤2 μg/10 ⁶ cells
PE conjugate	≤0.2 μg/10 ⁶ cells
APC conjugate	≤0.2 μg/10 ⁶ cells
SPRD conjugate	≤0.2 μg/10 ⁶ cells

Other Applications: Since applications vary, determine the optimum working dilution of the product that is appropriate for your specific needs.

HANDLING AND STORAGE

- The purified (UNLB) antibody is supplied as 0.5 mg of purified immunoglobulin in 1.0 mL of 100 mM borate buffered saline, pH 8.0. No preservatives or amine-containing buffer salts added.

- The fluorescein (FITC) conjugate is supplied as 0.5 mg in 1.0 mL of PBS/NaN₃.
- The biotin (BIOT) conjugate is supplied as 0.5 mg in 1.0 mL of PBS/NaN₃.
- The phycoerythrin (PE) conjugates are supplied as 0.1 mg in 1.0 mL or 0.2 mg in 2.0 mL of PBS/NaN₃ and a stabilizing agent.
- The allophycocyanin (APC) conjugate is supplied as 0.1 mg in 1.0 mL of PBS/NaN₃ and a stabilizing agent.
- The Spectral Red (SPRD) conjugate is supplied as 0.1 mg in 1.0 mL of PBS/NaN₃ and a stabilizing agent.
- Protect fluorochrome-conjugated forms from light. Do not freeze.
- Reagent is stable until the expiration date on the vial when stored at 2-8°C.

STATEMENT OF WARNINGS

1. Specimens, samples and all material coming in contact with them should be handled as if capable of transmitting infection and disposed of with proper precautions.
2. Never pipet by mouth and avoid contact of samples with skin and mucous membranes.
3. Do not use reagent beyond the expiration date on the vial label.
4. Minimize exposure of reagent to light during storage or incubation.
5. Avoid microbial contamination of reagent or erroneous results may occur.
6. Use Good Laboratory Practice (GLP) when handling this reagent.
7. Harmful if swallowed.
8. After contact with skin, wash immediately with plenty of water.
9. Contains sodium azide. Sodium azide under acidic conditions yields hydrazoic acid, an extremely toxic compound. Azide compounds should be flushed with running water while being discarded. These precautions are recommended to avoid deposits in metal piping in which explosive conditions can develop. If skin or eye contact occurs, immediately wash excessively with water.

TRADEMARKS

The Beckman Coulter logo is a trademark of Beckman Coulter, Inc.

Spectral Red is a trademark of Southern Biotechnology Associates, Inc.

Cy5 is a trademark of GE Healthcare, Inc.

For additional information or if damaged product is received, contact your local Beckman Coulter Representative.

Spectral Red is a PE/CyTM5 tandem conjugate. Cy5 is for non-commercial research use only, not for therapeutic or in vivo applications. Other use needs license from Amersham Biosciences Corp., under U.S. Patent Nos. 4,981,977 and 5,268,486 and other patents pending. This material (or portions of this material) is subject to proprietary rights of Amersham Biosciences Corp. and Carnegie Mellon University and made and sold under license from Amersham Biosciences Corp. This product is licensed for sale only for research. It is not licensed for any other use. There is no implied license hereunder for any commercial use. Commercial use shall include: 1) sale, lease, license or other transfer of the material or any material derived or produced from it 2) sale, lease, license or other grant of rights to use this material or any material derived or produced from it 3) use of this material to perform services for a fee for third parties. If you require a commercial license to use this material and do not have one, return this material, unopened to Beckman Coulter, Inc. 11800 SW 147 Ave. Miami, FL 33196, USA and any money paid for the material will be refunded.

REFERENCES

1. Gross JA, Callas E and Allison JP. 1992. Identification and distribution of the co-stimulatory receptor CD28 in the mouse. *J Immunol*, 149:380-388.
2. Giese T, Allison JP and Davidson WF. 1993. Functionally anergic lpr and gld B220+ T cell receptor (TCR)-alpha/beta+ double-negative T cells express CD28 and respond to co-stimulation with phorbol myristate acetate and antibodies to CD28 and the TCR. *J Immunol*, 151:597-609.
3. Clements JL, Winslow G, Donahue C, Cooper SM, Allison JP and Budd RC. 1993. Co-stimulation via CD28 induces activation of a refractory subset of MRL-lpr/lpr T lymphocytes. *Int Immunol*, 5:1451-1460.
4. Nandi D, Gross JA and Allison JP. 1994. CD28-mediated co-stimulation is necessary for optimal proliferation of murine NK cells. *J Immunol*, 152:3361-3369.

5. June CH, Bluestone JA, Nadler LM and Thompson CB. 1994. The B7 and CD28 receptor families. *Immunol Today*, 15:321-331.
6. Gelfanov V, Lai YG, Gelfanova V, Dong JY, Su JP and Liao NS. 1995. Differential requirement of CD28 co-stimulation for activation of murine CD8+ intestinal intraepithelial lymphocyte subsets and lymph node cells. *J Immunol*, 155:76-82.
7. Harding FA, McArthur JG, Gross JA, Raulet DH and Allison JP. 1992. CD28-mediated signalling co-stimulates murine T cells and prevents induction of anergy in T-cell clones. *Nature*, 356:607-609.
8. Bluestone JA. 1995. New perspectives of CD28-B7-mediated T cell co-stimulation. *Immunity*, 2:555-559.



Manufactured for:
Beckman Coulter, Inc.
4300 N. Harbor Blvd.
Fullerton, CA 92835
www.beckmancoulter.com

Printed in USA
Made in USA

© 2005 Beckman Coulter, Inc.
All Rights Reserved.

PN 733927-A